Neuroscience Letters, 93 (1988) 354–357
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Acworth, I.N., see During, M.J., 97
Agnit, L.F., see Bjelke, B., 139
Akaike, A., see Ujihara, H., 231
Alvarez-Bolado, G., see Cobas, A., 7
Amabile, G., see Strollo, F., 346
Anderson, C.R. and Howe, P.R.C., Is phenylethanolamine-N-methyltrasferase (PNMT) contained in rat hypothalamic neurons?, 164
Anderson, P.N., Turmaine, M. and Woodham, P., Schwann cells support extensive axonal growth into skeletal muscle implants in adult mouse brain, 127
Anderton, B.H., see Tagliavini, F., 191
Andreasen, M., Lambert, J.D.C. and Skovgaard Jensen, M., Direct demonstration of an N-methyl-D-aspartate receptor mediated component of excitatory synaptic transmission in area CA1 of the rat hippocampus, 61
Antonov, S.M. and Magazanik, L.G., Intense non-quantal release of glutamate in an insect neuromuscular junction, 204
Arvin, B., Neville, L.F. and Roberts, P.J., 2-Chloroadenosine prevents kainic acid-induced toxicity in rat striatum, 336
Aunis, D., see Scarfone, E., 13

Barragán, E. and Ferreyra-Moyano, H., Electrophysiological connections of neurons in ventral pallidal regions of the olfactory tubercle with the main olfactory bulb and piriform cortex, 214
Behzadi, G., see Wiklund, L., 158
Bicknell, R.J., Zhao, B.-G., Chaxman, C., Heavens, R.P. and Sirinathsinghji, D.J.S., Opioid inhibition of secretion from oxytocin and vasopressin nerve terminals following selective depletion of neurohypophysial catecholamines, 281
Bistritzer, T., Levin, P.A., Roeder, L.M. and Kapcala, L.P., Differential effects of adrenocorticotropin (1-24) on [^{3}H]2-deoxy-D-glucose uptake in cultured cells derived from different brain regions, 79
Bjelke, B., Fuxe, K. and Agnit, L.F., Survival of adrenohypophyseal homologous transplants in the rat striatum associated with prolactin-like immunoreactivity in the surrounding neuropil of the striatum, 139
Boden, P., see McBain, C.J., 341
Boegman, R.J., see Metcalf, R.H., 85

Carter, C., Noel, F. and Scatton, B., Sodium dependence of NMDA's effects on cyclic GMP production in immature rat cerebellar slices, 324
Chapman, C., see Bicknell, R.J., 281
Chase, T.N., see Mouradian, M.M., 225
Claps, A., see Torrealba, F., 186
Cobas, A., Alvarez-Bolado, G. and Fairén, A., Transient GABA-like immunoreactive axons in the corpus callosum of perinatal rats, 7
Cole, G.J., Schachner, M. and Fliesler, S.J., N-linked oligosaccharides are not required for neuron–neuron interactions mediated by neural cell adhesion molecule, 170
Colmers, W.F. and Williams, J.T., Pertussis toxin pretreatment discriminates between pre- and postsynaptic actions of baclofen in rat dorsal raphe nucleus in vitro, 300
Contreras, P.C., see Mouradian, M.M., 225

Datta, S.C., see Robertson, P.L., 107
Davrainville, J.-L., see Khayari, A., 56
Dememes, D., see Scarfone, E., 13
Dodson, A., see Tagliavini, F., 191
Dooley, D.J., Lickert, M., Lupp, A. and Osswald, H., Distribution of [^{125}I]ω-conotoxin GVIA and [^{3}H]isradipine binding sites in the central nervous system of rats of different ages, 318
Duan, X., see Ju, G., 19
During, M.J., Acworth, I.N. and Wurtman, R.J., Phenylalanine administration influences dopamine release in the rat's corpus striatum, 97

Engberg, G., see Sandberg, A.-C., 114

Fairén, A., see Cobas, A., 7
Fattapposta, F., see Strollo, F., 346
Feltz, A., see Stansfeld, C., 49
Ferreyra-Moyano, H., see Barragán, E., 214
Fliesler, S.J., see Cole, G.J., 170
Fritze, J., see Kornhuber, J., 312
Fu, Q.-G., see Sandkühler, J., 67
Fuller, R.W., see Gust, C.M., 307
Fuxe, K., see Bjelke, B., 139

Gallo, J.-M., see Tagliavini, F., 191

0304-3940/88/$ 03.50 © 1988 Elsevier Scientific Publishers Ireland Ltd.

Gharib, A., see Rogemont, C., 287
Goldstein, G.W., see Robertson, P.L., 107
Gust, C.M. and Fuller, R.W., Further characterization of tolerance to the corticosterone-elevating effect of pergolide in rats, 307

Hasegawa, Y., see Nakagawa, S., 32
Hashim, A., see Sershen, H., 270
Headley, P.M., see Wiklund, L., 158
Heavens, R.P., see Bicknell, R.J., 281
Heinsen, H., see Kornhuber, J., 312
Helmchen, C., see Sandkühler, J., 67
Hill, R.G., see McBain, C.J., 341
Howe, P.R.C., see Anderson, C.R., 164

Ishihara, M., see Satoh, M., 236

Jenner, P., see Nomoto, M., 275
Jiao, S.-S., see Lu, G.-W., 181
Jin, K.L., Shiotani, Y., Kawai, Y. and Kiyama, H., Immunohistochemical demonstration of tyrosine hydroxylase (TH)-positive but dopamine-hydroxylase (DBH)-negative neuron-like cells in the pineal gland of golden hamsters, 28
Johnston, G.A.R., see Nicholson, G.M., 330
Ju, G., Duan, X. and Zhang, X., Serotonin-like immunoreactive fibers in the medial dorsal accessory group and the anterior fornical nucleus of the magnocellular neurosecretory system in the rat, 19

Kalén, P., see Wiklund, L., 158
Kapcala, L.P., see Bistritzer, T., 79
Kato, M., Longitudinal myelotomy of lumbar spinal cord has little effect on coordinated locomotor activities of bilateral hindlimbs of the chronic cats, 259
Katsuki, H., see Satoh, M., 236
Kawai, Y., see Jin, K.L., 28
Khayari, A., Math, F. and Davrainville, J.-L., Evidence for an absence of K^+ spread in the glomerular layer of the rat olfactory bulb, 56
Kitao, Y., see Kudo, M., 176
Kiyama, H., see Jin, K.L., 28
Kobayashi, H., see Mochida, S., 247
Köller, H., see Siebler, M., 220
Kornhuber, J., Retz, W., Riederer, P., Heinsen, H. and Fritze, J., Effect of antemortem and postmortem factors on [^{3}H]glutamate binding in the human brain, 312
Kostowski, W., see Pucilowski, O., 96
Kovács, J., see Temesvári, P., 38
Kubozono, T., see Nakagawa, S., 32
Kudo, M., Nakamura, Y., Moriizumi, T., Tokuno, H. and Kitao, Y., Direct retinal projections to the lateroposterior thalamic nucleus (LP) in the mole, 176
Kumakura, K., see Ohara-Imaizumi, M., 294

Lajtha, A., see Sershen, H., 270
Lake, M., see Sandberg, A.-C., 114
Lakke, E.A.J.F. and Marini, E., A zonal pattern of neuroglial cells during the development of the intermediate lobe of the rat pituitary, 146
Lambert, J.D.C., see Andreasen, M., 61
Larsson, B.S., see Lindquist, N.G., 1
Levin, P.A., see Bistritzer, T., 79
Lickert, M., see Dooley, D.J., 318
Lindquist, N.G., Larsson. B.S. and Lydén-Sokolowski, A., Autoradiography of [^{14}C]paraquat or [^{14}C]diquat in frogs and mice: accumulation in neuromelanin, 1
Lobo, D.H. and Parnavelas, J.G., Ontogeny of neurotensin receptor binding sites in the rat visual cortex, 152
Lu, G.-W., Jiao, S.-S. and Zhang, G.-F., Morphological evidence for newly discovered double projection spinal neurons, 181
Ludwin, S.K., see Metcalf, R.H., 85
Lupp, A., see Dooley, D.J., 318
Lydén-Sokolowski, A., see Lindquist, N.G., 1

MacDonald, J.F., see Mody, I., 73
Magari, S., see Sakanaka, M., 23
Magazanik, L.G., see Antonov, S.M., 204
Marini, E., see Lakke, E.A.J.F., 146
Markovac, J., see Robertson, P.L., 107
Marsden, C.D., see Nomoto, M., 275
Martini, M., see Rossi, M.L., 44
Math, F., see Khayari, A., 56
Matsumoto, N., Sato, T., Sawano, H., Tochinai, A. and Suzuki, T.A., Characteristics of tooth pulp-driven neurons in the posterior group of the cat thalamus, 253
McBain, C.J., Boden, P. and Hill, R.G., The kainate/quisqualate receptor antagonist, CNQX, blocks the fast component of spontaneous epileptiform activity in organotypic cultures of rat hippocampus, 341
McGarrick, J.D., see Thexton, A.J., 242
Mengod, G., see Savasta, M., 132
Metcalf, R.H., Boegman, R.J., Riopelle, R.J. and Ludwin, S.K., The release of endogenous acetylcholine from the medial septum/diagonal band of rat brain, 85
Meyer, D.L., see Weiss, 0., 209

Miyakawa, Y., see Ohara-Imaizumi, M., 294
Mochida, S. and Kobayashi, H., GTP-binding proteins mediate the M_2-muscarinic effect on the action potential in isolated sympathetic neurons of rabbits, 247
Mody, I., Salter, M.W. and MacDonald, J.F., Requirement of NMDA receptor/channels for intracellular high-energy phosphates and the extent of intraneuronal calcium buffering in cultured mouse hippocampal neurons, 73
Monahan, J.B., see Mouradian, M.M., 225
More, M., see Strollo, F., 346
Moriizumi, T., see Kudo, M., 176
Mouradian, M.M., Contreras, P.C., Monahan, J.B. and Chase, T.N., [^{3}H]MK-801 binding in Alzheimer's disease, 225
Müller, H.W., see Siebler, M., 220

Nakagawa, S., Hasegawa, Y., Kubozono, T. and Takuni, K., Substance P-like immunoreactive retinal terminals found in two retinoreceipient areas of the Japanese monkey, 32
Nakamura, Y., see Kudo, M., 176
Neville, L.F., see Arvin, B., 336
Nicholson, G.M., Spence, I. and Johnston, G.A.R., Depolarizing actions of convulsant barbiturates on isolated rat dorsal root ganglion cells, 330
Nicolopoulos, L.S., see Wiklund, L., 158
Noel, F., see Carter, C., 324
Nomoto, M., Jenner, P. and Marsden, C.D., The D_1 agonist SKF 38393 inhibits the antiparkinsonian activity of the D_2 agonist LY 171555 in the MPTP-treated marmoset, 275

Ohara-Imaizumi, M., Miyakawa, Y. and Kumakura, K., Pertussis toxin attenuates clonidine inhibition of catecholamine release in adrenal chromaffin cells, 294
Olley, J.E., see Tiong, G.K.L., 101
Osswald, H., see Dooley, D.J., 318

Pacheco, H., see Rogemont, C., 287
Palacios, J.M., see Savasta, M., 132
Parnavelas, J.G., see Lobo, D.H., 152
Parsons, C.G., see Wiklund, L., 158
Pasingham, R.E., see Thaler, D.E., 264
Perrin, D., see Scarfone, E., 13
Power, D.M., see Tagliavini, F., 191
Pucilowski, O. and Kostowski, W., Diltiazem suppresses apomorphine-induced fighting and pro-aggressive effect of withdrawal from chronic ethanol or haloperidol in rats, 96

Retz, W., see Kornhuber, J., 312
Riederer, P., see Kornhuber, J., 312
Riondino, G., see Strollo, F., 346
Riopelle, R.J., see Metcalf, R.H., 85
Roberts, P.J., see Arvin, B., 336
Robertson, P.L., Markovac, J., Datta, S.C. and Goldstein, G.W., Transforming growth factor beta stimulates phosphoinositol metabolism and translocation of protein kinase C in cultured astrocytes, 107
Roeder, L.M., see Bistritzer, T., 79
Rogemont, C., Sarda, N., Gharib, A. and Pacheco, H., Changes in the rat sleep-wake cycle produced by D,L-(1-naphthyl)alanine, a tryptophan analog, 287
Rolls, E.T., see Thaler, D.E., 264
Rossi, M.L. and Martini, M., High perilymphatic potassium and rubidium levels modulate the sensory resting discharge in the isolated frog labyrinth, 44

Sakanaka, M., Magari, S. and Shibasaki, T., Fine structures of nerve fibers with corticotropin-releasing factor-like immunoreactivity in the rat lateral septum, 23
Salter, M.W., see Mody, I., 73
Sandberg, A.-C., Engberg, G., Lake, M., Von Holst, H. and Sara, V.R., The expression of insulin-like growth factor I and insulin-like growth factor II genes in the human fetal and adult brain and in glioma, 114
Sandkühler, J., Helmchen, C., Fu, Q.-G. and Zimmermann, M., Inhibition of spinal nociceptive neurons by excitation of cell bodies or fibers of passage at various brainstem sites in the cat, 67
Sans, A., see Scarfone, E., 13
Sara, V.R., see Sandberg, A.-C., 114
Sarda, N., see Rogemont, C., 287
Sasa, M., see Ujihara, H., 231
Sato, T., see Matsumoto, N., 253
Satoh, M., Ishihara, M. and Katsuki, H., Different susceptibilities of long-term potentiations in CA3 and CA1 regions of guinea pig hippocampal slices to nootropic drugs, 236
Savasta, M., palacios, J.M. and Mengod, G., Regional localization of the mRNA coding for the neuropeptide cholecystokinin in the rat brain studied by in situ hybridization, 132
Sawano, H., see Matsumoto, N., 253
Scarfone, E., Dememes, D., Perrin, D., Aunis, D. and Sans, A., α-Fodrin (brain spectrin) immunocytochemical localization in rat vestibular hair cells, 13

Scatton, B., see Carter, C., 324
Schachner, M., see Cole, G.J., 170
Schmalenbach, C., see Siebler, M., 220
Sershen, H., Hashim, A., Wiener, H.L. and Lajtha, A., Effect of chronic oral nicotine on dopaminergic function in the MPTP-treated mouse, 270
Shibasaki, T., see Sakanaka, M., 23
Shiotani, Y., see Jin, K.L., 28
Siebler, M., Köller, H., Schmalenbach, C. and Müller, H.W., GABA activated chloride currents in cultured rat hippocampal and septal region neurons can be inhibited by curare and atropine, 220
Sirinathsinghji, D.J.S., see Bicknell, R.J., 281
Skovgaard Jensen, M., see Andreasen, M., 61
Spence, I., see Nicholson, G.M., 330
Stansfeld, C. and Feltz, A., Dendrotoxin-sensitive K^+ channels in dorsal root ganglion cells, 49
Stone, T.W., see Thexton, A.J., 242
Strollo, F., Amabile, G., Fattapposta, F., Strollo, G., More, M. and Riondino, G., Thyrotropin-releasing hormone enhances event-related brain potentials and growth hormone release in man, 346
Strollo, G., see Strollo, F., 346
Suzuki, T.A., see Matsumoto, N., 253
Swash, M., see Tagliavini, F., 191

Tagliavini, F., Anderton, B.H., Dodson, A., Gallo, J.-M., Swash, M. and Power, D.M., Preamyloid deposits in the cerebral cortex of patients with Alzheimer's disease and non-demented individuals, 191
Takaori, S., see Ujihara, H., 231
Takuni, K., see Nakagawa, S., 32
Temesvári, P. and Kovács, J., Selective opening of the blood–brain barrier in newborn piglets with experimental pneumothorax, 38
Thaler, D.E., Rolls, E.T. and Pasingham, R.E., Neuronal activitity of the supplementary motor area (SMA) during internally and externally triggered wrist movements, 264
Thexton, A.J., McGarrick, J.D. and Stone, T.W., Rhythmic digastric activity in the naloxone-treated decerebrate rabbit pup, 242
Tiong, G.K.L. and Olley, J.E., Effects of exposure in utero to methadone and buprenorphine on enkephalin levels in the developing rat brain, 101
Tochinai, A., see Matsumoto, N., 253
Tokuno, H., see Kudo, M., 176
Torrealba, F. and Claps, A., The vagal connection of the carotid sinus, 186
Turmaine, M., see Anderson, P.N., 127

Ujihara, H., Akaike, A., Sasa, M. and Takaori, S., Electrophysiological evidence for cholinoceptive neurons in the medial vestibular nucleus: studies on rat brain stem in vitro, 231

Von Holst, H., see Sandberg, A.-C., 114

Wathey, J.C. and Wullimann, M.F., A double-label study of efferent projections from the Edinger-Westphal nucleus in goldfish and kelp bass, 121
Weiss, O. and Meyer, D.L., Odor stimuli modulate retinal excitability in fish, 209
West, D.C., see Wiklund, L., 158
Wiener, H.L., see Sershen, H., 270
Wiklund, L., Behzadi, G., Kalén, P., Headley, P.M., Nicolopoulos, L.S., Parsons, C.G. and West, D.C., Autoradiographic and electrophysiological evidence for excitatory amino acid transmission in the periaqueductal gray projection to nucleus raphe magnus in the rat, 158
Williams, J.T., see Colmers, W.F., 300
Woodham, P., see Anderson, P.N., 127
Wullimann, M.F., see Wathey, J.C., 121
Wurtman, R.J., see During, M.J., 97

Zhang, G.-F., see Lu, G.-W., 181
Zhang, X., see Ju, G., 19
Zhao, B.-G., see Bicknell, R.J., 281
Zimmermann, M., see Sandkühler, J., 67